

A counterexample against the Vlasov equation

C. Y. Chen

Department of Physics, Beijing University of Aeronautics
and Astronautics, Beijing, 100083, P.R.China *

April 21, 2009

Abstract

A simple counterexample against the Vlasov equation is put forward, in which a magnetized plasma is perturbed by an electromagnetic standing wave.

In today's physics, the Vlasov equation serves as the very foundation for a great number of formulations and calculations in fields ranging from star physics to controlled fusion.

The equation states that the distribution function $f(t, \mathbf{r}, \mathbf{v})$ of a collection of moving charged particles (called a plasma in the usual physics language) satisfies[1, 2]

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{r}} + \frac{q}{m}(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \frac{\partial f}{\partial \mathbf{v}} = 0, \quad (1)$$

where $(\mathbf{E}, \mathbf{B})$ represents the electromagnetic field experienced by the particles of charge q and mass m at the position $\mathbf{r}$ and with the velocity $\mathbf{v}$. It is sometimes called the collisionless Boltzmann equation in statistical mechanics due to the fact that the equation takes almost the same form of the ordinary Boltzmann equation with the collision term neglected. In an alternative form, Eq. (1) is written as[1, 3]

$$\left. \frac{df[t, \mathbf{r}(t), \mathbf{v}(t)]}{dt} \right|_{\text{path}} = 0, \quad (2)$$

*Email: cychen@buaa.edu.cn

where the subscripts imply that the derivative is performed along a particle's path. This expression can be interpreted as the path-invariance of the distribution function in the six-dimensional phase space spanned by the position $\mathbf{r} = (x, y, z)$ and velocity $\mathbf{v} = (v_x, v_y, v_z)$. Recognized as a good statistic approximation by this community, Eq. (1), or (2), is very often used to investigate the dynamical behavior of dilute plasma.

Being exposed by the fact that the regular Boltzmann equation suffers substantial difficulties[4, 5], the author has somehow been compelled to find out in what situations the Vlasov equation can be falsified. It turns out that a number of counterexamples against the Vlasov equation can be constructed in plain manners. In this paper, we shall focus ourselves on one of them, in which the six-dimensional physics is reduced to a one-dimensional one and can be understood with conceptual clarity.

Let's start with a plasma having a strong constant magnetic field B_0 along the z -direction. Suppose that its distribution function $f \equiv f(t, \mathbf{r}, \mathbf{v})$ is initially uniform in the position space, and that it is initially independent of velocity when the speeds of the concerned particles are below a certain limit v_0 . Namely, we have, with $v < v_0$ and at $t = 0$,

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = \frac{\partial f}{\partial z} = \frac{\partial f}{\partial v_x} = \frac{\partial f}{\partial v_y} = \frac{\partial f}{\partial v_z} = 0; \quad (3)$$

or, in another form,

$$f(t, \mathbf{r}, \mathbf{v})|_{t=0} = f_c, \quad (4)$$

where f_c represents a constant. For further simplicity, we also assume that the particles whose speeds are beyond v_0 occupy only a small proportion of the plasma and can be disregarded in our consideration.

Adopting all the initial conditions mentioned above, we see that no matter what kind of electromagnetic field is applied, Eq. (1) leads us to nothing but

$$\left. \frac{\partial f}{\partial t} \right|_{t=0} = 0, \quad (5)$$

which means that the distribution function of the system does not change at the beginning. More than that, Eq. (2) informs us that if an observer adheres to any moving particle, he/she finds that f is equal to f_c constantly, irrespective of the position and velocity, which literally infers that the distribution function remains unchanged at the beginning and the later time; or, in terms of macroscopic quantities there is no density change and no current everywhere once for all (readers may also refer to Chapter 8 of Ref. 1, in which the linearized Vlasov equation for uniformly magnetized plasmas yields the same conclusion).

However, physical inspection tells us a very different story: applying many types of electromagnetic fields will change the plasma significantly. For this to be seen in an intuitive and heuristic manner, let the magnetized plasma be perturbed by an electromagnetic standing wave in the form:

$$E_{1y} = \frac{a}{k_x} \cos(\omega t) \sin(k_x x), \quad B_{1z} = -\frac{a}{\omega} \sin(\omega t) \cos(k_x x), \quad (6)$$

where a is a small constant. Note that this perturbative wave obeys Maxwell's equation $\nabla \times \mathbf{E} = -\partial \mathbf{B} / \partial t$.

As well known in physics, while all particles in the plasma make gyration due to the existence of B_0 , the gyro-center of every particle (guiding center) involves two types of relatively slow drifts due to the existence of E_{1y} and B_{1z} . The first slow drift is the $\mathbf{E} \times \mathbf{B}$ drift along the x -direction:

$$\bar{v}_x = \frac{E_{1y}}{B_0} = \frac{a}{k_x B_0} \cos(\omega t) \sin(k_x x), \quad (7)$$

and the second slow drift the grad-B drift along the y -direction

$$\bar{v}_y = \frac{v_{\perp} \rho}{2B_0} \cdot \frac{\partial B_{1z}}{\partial x} = \frac{a k_x v_{\perp} \rho}{2\omega B_0} \sin(\omega t) \sin(k_x x). \quad (8)$$

Equations (7) and (8) hold when ω is much smaller than the gyro-frequency and the perturbed fields are relatively weak.

If observing the plasma in terms of measuring velocity and speed, two things can be found. The first is that expression (7) apparently represents an average velocity along the x -direction (forming a macroscopic current if the plasma is a single-component one). The second is that there is a speed change related to every particle due to the combined effect of the grad-B drift and the electric field E_{1y} although the change, of a^2 -order in magnitude, is relatively unimportant in a perturbative approach.

If the density change is of concern, something against the standard theory will also surface. While the grad-B drift expressed by (8) is uniform along the y -direction, causing no density change anywhere, the $\mathbf{E} \times \mathbf{B}$ drift expressed by (7) is nonuniform along the x -direction and must result in a density change along the x -direction. By assuming that the gyro-radii of all the concerned particles are relatively small (much smaller than $2\pi/k_x$) and noticing the fact that the $\mathbf{E} \times \mathbf{B}$ drift is velocity-independent, the density change can be evaluated with ease. With help of the linearized continuity equation of ordinary fluid, we find that

$$\frac{\partial n_1}{\partial t} = -n_0 \frac{\partial \bar{v}_x}{\partial x} = -\frac{n_0 a}{B_0} \cos(\omega t) \cos(k_x x), \quad (9)$$

where n_0 is the initial density of the plasma. Fig. 1 demonstrates that particles in this plasma, regardless of their velocities, converge and diverge in the pattern revealed by (9).




Fig. 1: The $\mathbf{E} \times \mathbf{B}$ drift of gyrating particles along the x -direction caused by the electromagnetic standing wave.

It is at this stage intriguing and essential to look at what happens in the phase space spanned by x, y, z, v_x, v_y, v_z . It should be pointed out that this six-dimensional space is a truly abstract and weird one, serving as a great source of all confusions and mistakes. Investigating the change of f in the phase space challenges our conceptual and experimental wisdom though it is surely in the domain of classical mechanics. Referring to Fig. 1, let's imagine that there is a small virtual box adhering to a moving particle, represented by $\Delta x \Delta y \Delta z$ symbolically, and imagine that we are able to count the number of the inside particles whose velocities are within a small range, represented by $\Delta v_x \Delta v_y \Delta v_z$ symbolically. What will be seen by us when the box $\Delta x \Delta y \Delta z$ drifts leftwards and rightwards in the circumstances? At the two ending points, the density in the box $\Delta x \Delta y \Delta z$ will, as revealed above, change from the lowest to the highest (or vice versa) while the velocities of the particles inside the box keep almost unchanged (noticing that the drift velocity at the ending points are zero and the speed changes of the particles are of second order). All this means that f in the phase volume element $\Delta x \Delta y \Delta z \Delta v_x \Delta v_y \Delta v_z$ is not invariant along a particle's path.

Many theoretical and practical questions then arise. If the Vlasov equation is indeed unsound, how come there exist so many academical proofs showing otherwise? If the standard theory fails to yield good predictions about density's changes and drift currents, why haven't present computa-

tional simulations revealed any discrepancy between the theory and reality? In what way can we correctly formulate the behavior of plasma? Extensive effort has been made by the author to answer these questions and to find out new things[5, 6], but the greatest challenge was, and still is, to convince the mainstream of this community to have open mind on related issues.

The author thanks professors Oliver Penrose and Keying Guan for stimulating discussions.

References

- [1] N.A. Krall and A.W. Trivelpiece, *Principle of Plasma Physics*, (McGraw-Hill Book Company, 1973).
- [2] S. Ichimaru, *Basic Principle of Plasma Physics*, (W. A. Benjamin, 1973).
- [3] F. Reif, *Fundamentals of Statistical and Thermal Physics*, (McGraw-Hill Book Company, 1965, 1987, 1988 in English and German).
- [4] C.Y. Chen, *Mathematical Investigation of the Boltzmann Collisional Operator*, II Nuovo Cimento B **V117B**, 177-181 (2002).
- [5] C.Y. Chen, *A new uncertainty principle*, arxiv.org: 0812.4343, 2008.
- [6] C.Y. Chen, *Perturbation Methods and Statistical Theories*, in English, (International Academic Publishers, Beijing, 1999); and cond-mat/0608712, 0504497, 0412396; physics/0312043, 0311120, 0305006, 0010015, 0006033, 0006009, 9908062.